

PATIENT INFORMATION

Propafenone Hydrochloride Extended-Release Capsules, USP (proe" pa fee' none hye" droe klor' ide)

What are propafenone hydrochloride extended-release capsules?

Propafenone hydrochloride extended-release capsules are a prescription medicine that is used:

- in certain people who have a heart rhythm disorder called atrial fibrillation (AF)
- to increase the amount of time between having symptoms of AF

It is not known if propafenone hydrochloride extended-release capsules are safe and effective in children.

Who should not take propafenone hydrochloride extended-release capsules?

Do not take propafenone hydrochloride extended-release capsules if you have:

- heart failure (weak heart)
- had a recent heart attack
- a heart condition called Brugada Syndrome
- a heart rate that is too slow, and you do not have a pacemaker
- very low blood pressure
- certain breathing problems that make you short of breath or wheeze
- certain abnormal body salt (electrolyte) levels in your blood

Talk to your doctor before taking propafenone hydrochloride extended-release capsules if you think you have any of the conditions listed above.

What should I tell my doctor before taking propafenone hydrochloride extended-release capsules?

Before you take propafenone hydrochloride extended-release capsules, tell your doctor if you:

- have liver or kidney problems
- have breathing problems
- have symptoms including diarrhea, sweating, vomiting, or loss of appetite or thirst that are severe. These symptoms may be a sign of abnormal electrolyte levels in your blood.
- have myasthenia gravis
- have lupus erythematosus
- have been told you have or had an abnormal blood test called Antinuclear Antibody Test or ANA Test
- are pregnant or plan to become pregnant
- are breastfeeding or plan to breastfeed. Propafenone hydrochloride extended-release capsules can pass into your milk. You and your doctor should discuss the best way to feed your baby during this time.
- have any other medical conditions

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Propafenone hydrochloride extended-release capsules and certain other medicines can affect each other and cause serious side effects. Propafenone hydrochloride extended-release capsules may affect the way other medicines work, and other medicines may affect how propafenone hydrochloride extended-release capsules work.

Especially tell your doctor if you take:

- amiodarone or other medicines for your abnormal heart beats
- an antidepressant medicine
- a medicine to treat anxiety
- ritonavir (for example, KALETRA, NORVIR) or saquinavir (for example, INVIRASE)
- an antibiotic medicine
- ketoconazole (for example, NIZORAL)
- digoxin (LANOXIN)
- warfarin sodium (for example, COUMADIN, JANTOVEN)

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

How should I take propafenone hydrochloride extended-release capsules?

- Take propafenone hydrochloride extended-release capsules exactly as prescribed. Your doctor will tell you how many capsules to take and how often to take them.
- To help reduce the chance of certain side effects, your doctor may start you with a low dose of propafenone hydrochloride extended-release capsules, and then slowly increase the dose.
- Do not open or crush the capsule.
- You may take propafenone hydrochloride extended-release capsules with or without food.
- You should not drink grapefruit juice during treatment with propafenone hydrochloride extended-release capsules.
- If you miss a dose of propafenone hydrochloride extended-release capsules, take your next dose at the usual time. Do not take 2 doses at the same time.
- If you take too much propafenone hydrochloride extended-release capsules, call your doctor or go to the nearest hospital emergency room right away.
- Call your doctor if your heart problems get worse.

What are possible side effects of propafenone hydrochloride extended-release capsules?

Propafenone hydrochloride extended-release capsules can cause serious side effects including:

- **New or worsened abnormal heart beats, that can cause sudden death or be life-threatening.** Your doctor may do an electrocardiogram (ECG or EKG) before and during treatment to check your heart for these problems.
- **New or worsened heart failure. Tell your doctor about any changes in your heart symptoms, including:**
 - any new or increased swelling in your arms or legs
 - trouble breathing
 - sudden weight gain

- **Effects on pacemaker function.** Propafenone hydrochloride extended-release capsules may affect how an implanted pacemaker or defibrillator works. Your doctor should check how your pacemaker or defibrillator is working during and after treatment with propafenone hydrochloride extended-release capsules. They may need to be re-programmed.
- **Very low white blood cell levels in your blood (agranulocytosis).** Your bone marrow may not produce enough of a certain type of white blood cells called neutrophils. If this happens, you are more likely to get infections. Tell your doctor right away if you have any of these symptoms, especially during the first 3 months of treatment:
 - fever
 - sore throat
 - chills
- **Worsening of myasthenia gravis in people who already have this condition.** Tell your doctor about any change in your symptoms.
- **Propafenone hydrochloride extended-release capsules may cause lower sperm counts in men.** This could affect the ability to father a child. Talk to your doctor if this is a concern for you.

Common side effects of propafenone hydrochloride extended-release capsules include:

- dizziness
- fast or irregular heart beats
- chest pain
- trouble breathing
- taste changes
- nausea
- tiredness
- feeling anxious
- constipation
- upper respiratory infection or flu
- swelling

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of propafenone hydrochloride extended-release capsules. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store propafenone hydrochloride extended-release capsules?

- Store propafenone hydrochloride extended-release capsules at 20°C to 25°C (68°F to 77°F).
- Keep the bottle tightly closed.

Keep propafenone hydrochloride extended-release capsules and all medicines out of the reach of children.

General information about propafenone hydrochloride extended-release capsules

Medicines are sometimes prescribed for conditions other than those described in patient information leaflets. Do not use propafenone hydrochloride extended-release capsules for a condition for which it was not prescribed by your doctor. Do not give propafenone hydrochloride extended-release capsules to other people, even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about propafenone hydrochloride extended-release capsules. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about propafenone hydrochloride extended-release capsules that is written for healthcare professionals. For more information about propafenone hydrochloride extended-release capsules, call 1-844-518-2989.

What are the ingredients in propafenone hydrochloride extended-release capsules?

Active Ingredient: Propafenone hydrochloride

Inactive Ingredients: gelatin, hypromellose, magnesium stearate, potassium hydroxide, propylene glycol, red iron oxide, shellac, sodium lauryl sulfate, strong ammonia solution and titanium dioxide.

The brands listed are trademarks of their respective owners.

Manufactured for:

TWi Pharmaceuticals USA, Inc.
Paramus, NJ 07652

Manufactured by:

TWI

TWi Pharmaceuticals, Inc.
Taoyuan City, 32063, Taiwan

Revised: 06/2020

REVIEWED

By Katy Saucedo at 2:06 pm, Sep 07, 2021

APPROVED

By mdlrosa at 12:35 pm, Sep 10, 2021